



Age Defense Eye Cream Gel with Hyasol-BT and REGU®-Age

<u>Phase A</u>	<u>%W/W</u>
Ethylhexyl Isononanoate (Pelemol 89/Phoenix Chemicals)	5.00
Isohexadecane (Arlamol HD/Uniqema)	2.00
Diethylhexyl Carbonate (Tegosoft DEC/Evonik)	5.00
Dimethicone (DC200fluid/200cts/Dow Corning)	1.00
 <u>Phase B</u>	 <u>%W/W</u>
Ammonium Acryloyldimethyltaurate & VP Copolymer (Arisoflex AVC/Clariant)	1.20
 <u>Phase C</u>	 <u>%W/W</u>
Deionized Water	72.10
Glycerin (Rita Glycerin USP/Rita)	5.00
Sodium Hydroxide (10% solution)	q.s. to pH 5.5—8.0
 <u>Phase D</u>	 <u>%W/W</u>
¹ REGU-AGE	5.00
¹ HYASOL-BT	3.00
² DIOCIDE	0.70

¹Manufactured by DSM Nutritional Products Pentapharm Branch, North American Distributor: Centerchem, Inc.

²Centerchem, Inc.

Procedure:

1. Mix Phase A together. Add Phase B to Phase A, mix until homogeneous.
2. Mix together Phase C ingredients. Adjust pH to 5.5—8.0, if necessary.
3. Add Phase D ingredients to Phase C.
4. Add Phase A/B to Phase C/D. Mix to obtain a homogenous cream gel. Discontinue.

Ref. D100-082

This information and all further technical advice is based on our present knowledge and experience. However, it implies no liability, or other legal responsibility on our part, including with regard to existing third party intellectual property rights, especially patent rights. In particular, no warranty, whether express or implied, or guarantee of product properties in the legal sense is intended or implied. We reserve the right to make any changes according to technological progress or further developments.

