



Moisturizing Hair Gel *with* PENTAVITIN®

<u>Phase A</u>	<u>% W/W</u>
Deionized Water	65.50
Carbomer (Ultrez 10/Lubrizol)	0.40
Glycerin	20.00
<u>Phase B</u>	<u>% W/W</u>
¹ HISPAGEL 200NS	10.00
<u>Phase C</u>	<u>% W/W</u>
² PENTAVITIN	3.00
<u>Phase D</u>	<u>% W/W</u>
Triethanolamine 99%	0.40
<u>Phase E</u>	<u>% W/W</u>
³ DIOCIDE	0.70

¹Manufactured by Cognis Iberia S.L., Exclusive North American Distributor: Centerchem, Inc.

²Manufactured by DSM Nutritional Products Pentapharm Branch, North American Distributor: Centerchem, Inc.

³Centerchem, Inc.

Procedure:

1. Sprinkle carbomer in Phase A water with mixing. Once dispersed, add glycerin.
2. Add Phase B, mix until uniform.
3. Add Phase C, mix until uniform.
4. Add Phase D, mix until uniform.
5. Add Phase E, mix until uniform.

Ref. F100-104

This information and all further technical advice is based on our present knowledge and experience. However, it implies no liability, or other legal responsibility on our part, including with regard to existing third party intellectual property rights, especially patent rights. In particular, no warranty, whether express or implied, or guarantee of product properties in the legal sense is intended or implied. We reserve the right to make any changes according to technological progress or further developments.

